arXiv:1603.01850v1 [math.AC] 6 Mar 2016

TORIC RINGS AND IDEALS OF STABLE SET POLYTOPES

KAZUNORI MATSUDA, HIDEFUMI OHSUGI AND KAZUKI SHIBATA

Abstract. In this paper, we discuss the normality of the toric rings of stable set polytopes, and the set of generators and Gröbner bases of toric ideals of stable set polytopes by using the results on that of edge polytopes of finite nonsimple graphs. In particular, for a graph of stability number two, we give a graph theoretical characterization of the set of generators of the toric ideal of the stable set polytope, and a criterion to check whether the toric ring of the stable set polytope is normal or not. One of the application of the results is an infinite family of stable set polytopes whose toric ideal is generated by quadratic binomials and has no quadratic Gröbner bases.

Introduction

Let $\mathcal{P} \subset \mathbb{R}^n$ be an integral convex polytope, i.e., a convex polytope each of whose vertices has integer coordinates, and let $\mathcal{P} \cap \mathbb{Z}^n = \{\mathbf{a}_1, \ldots, \mathbf{a}_m\}$. Let $K[X, X^{-1}, t] = K[x_1, x_1^{-1}, \ldots, x_n, x_n^{-1}, t]$ be the Laurent polynomial ring in $n+1$ variables over a field K. Given an integer vector $\mathbf{a} = (a_1, \ldots, a_n) \in \mathbb{Z}^n$, we set $X^{\mathbf{a}}t = x_1^{a_1} \cdots x_n^{a_n} t \in K[X, X^{-1}, t]$. Then, the *toric ring* of $\mathcal{P}$ is the subalgebra $K[\mathcal{P}]$ of $K[X, X^{-1}, t]$ generated by $\{X^{\mathbf{a}_1}t, \ldots, X^{\mathbf{a}_m}t\}$ over K. We regard $K[\mathcal{P}]$ as a homogeneous algebra by setting each $\deg X^{\mathbf{a}_i}t = 1$. The *toric ideal* $I_{\mathcal{P}}$ of $\mathcal{P}$ is the kernel of a surjective homomorphism $\pi : K[y_1, \ldots, y_m] \to K[\mathcal{P}]$ defined by $\pi(y_i) = X^{\mathbf{a}_i}t$ for $1 \leq i \leq m$. It is known that $I_{\mathcal{P}}$ is generated by homogeneous binomials and reduced Gröbner bases of $I_{\mathcal{P}}$ consist of homogeneous binomials. See, e.g., [20]. The following properties on an integral convex polytope $\mathcal{P}$ have been investigated by many papers on commutative algebra and combinatorics:

(i) $\mathcal{P}$ is unimodular, i.e., any triangulation of $\mathcal{P}$ is unimodular
(The initial ideal of $I_{\mathcal{P}}$ is generated by squarefree monomials with respect to any monomial order);
(ii) $\mathcal{P}$ is compressed, i.e., any "pulling" triangulation is unimodular
(The initial ideal of $I_{\mathcal{P}}$ is generated by squarefree monomials with respect to any reverse lexicographic order);
(iii) $\mathcal{P}$ has a regular unimodular triangulation
(There exists a monomial order such that the initial ideal of $I_{\mathcal{P}}$ is generated by squarefree monomials);
(iv) $\mathcal{P}$ has a unimodular triangulation;
(v) $\mathcal{P}$ has a unimodular covering;
(vi) $\mathcal{P}$ is normal, i.e., $K[\mathcal{P}]$ is normal.

Key words and phrases. toric ideals, Gröbner bases, graphs, stable set polytopes.

The hierarchy (i) $\Rightarrow$ (ii) $\Rightarrow$ (iii) $\Rightarrow$ (iv) $\Rightarrow$ (v) $\Rightarrow$ (vi) is known. See, e.g., [20] for details. However, the converse of each of the five implications is false. On the other hand, the following properties on $I_{\mathcal{P}}$ are studied by many authors:

(a) $I_{\mathcal{P}}$ has a quadratic Gröbner bases;
(b) $K[\mathcal{P}]$ is Koszul algebra;
(c) $I_{\mathcal{P}}$ is generated by quadratic binomials.

The hierarchy (a) $\Rightarrow$ (b) $\Rightarrow$ (c) is known. However, the converse of each of the two implications is false. See, e.g., [13, Examples 2.1 and 2.2] and [21].

The purpose of this paper is to study such properties of toric rings and ideals of stable set polytopes of simple graphs. Let G be a finite simple graph on the vertex set $[n] = \{1, 2, \ldots, n\}$ and let $E(G)$ denote the set of edges of G. Given a subset $W \subset [n]$, we associate the $(0,1)$-vector $\rho(W) = \sum_{j \in W} \mathbf{e}_j \in \mathbb{R}^n$. Here, $\mathbf{e}_i$ is the ith unit coordinate vector of $\mathbb{R}^n$. In particular, $\rho(\emptyset)$ is the origin of $\mathbb{R}^n$. A subset W is called *stable* if $\{i, j\} \notin E(G)$ for all $i, j \in W$ with $i \neq j$. Note that the empty set and each single-element subset of $[n]$ are stable. Let $S(G)$ denote the set of all stable sets of G. The *stable set polytope* (*independent set polytope*) of a simple graph G is the $(0,1)$-polytope $\mathcal{Q}_G \subset \mathbb{R}^n$ which is the convex full of $\{\rho(W) : W \in S(G)\}$. Stable set polytopes are very important in many areas, e.g., optimization theory. On the other hand, several results are known for the toric ring $K[\mathcal{Q}_G]$ and ideals $I_{\mathcal{Q}_G}$ of the stable set polytope $\mathcal{Q}_G$ of a simple graph G.

(1) For a graph G, the stable set polytope $\mathcal{Q}_G$ is compressed if and only if G is perfect ([5, 15, 22]).
(2) For a perfect graph G, the toric ring $K[\mathcal{Q}_G]$ is Gorenstein if and only if all maximal cliques of G have the same cardinality ([16]).
(3) For a graph G, $K[\mathcal{Q}_G]$ is strongly Koszul if and only if G is trivially perfect ([7, Theorem 5.1]).
(4) Let $G(P)$ be a comparability graph of a poset P. Then, $\mathcal{Q}_{G(P)}$ is called a *chain polytope* of P. It is known that the toric ideals of a chain polytope has a squarefree quadratic initial ideal (see [6, Corollary 3.1]). For example, if a graph G is bipartite, then there exists a poset P such that $G = G(P)$.
(5) Suppose that a graph G on the vertex set $[n]$ is an almost bipartite graph, i.e., there exists a vertex v such that the induced subgraph of G on the vertex set $[n] \setminus \{v\}$ is bipartite. Then, $I_{\mathcal{Q}_G}$ has a squarefree quadratic initial ideal ([2, Theorem 8.1]). For example, any cycle is almost bipartite.
(6) Let G be the complement of even cycle of length $2k$. Then, the maximum degree of a minimal set of binomial generators of $I_{\mathcal{Q}_G}$ is equal to k ([2, Theorem 7.4]).

In the present paper, we study the normality of the toric rings of stable set polytopes, and the set of generators and Gröbner bases of toric ideals of stable set polytopes by using the results on that of edge polytopes of finite (nonsimple) graphs. Here, the *edge polytope* $\mathcal{P}_G \subset \mathbb{R}^n$ of a graph G allowing loops and having no multiple edges is the convex full of

$$\{\mathbf{e}_i + \mathbf{e}_j : \{i, j\} \text{ is an edge of } G\} \cup \{2\mathbf{e}_i : \text{ there is a loop at } i \text{ in } G\}.$$

This paper is organized as follows. In Section 1, fundamental properties of $K[\mathcal{Q}_G]$ and $I_{\mathcal{Q}_G}$ are studied. In particular, the relationship between the stable set polytopes and the edge polytopes are given (Lemma 1.7). In addition, it is shown that $\mathcal{Q}_G$ is unimodular if and only if the complement of G is bipartite (Proposition 1.8). We also point out that, by the results in [2], it is easy to see that $I_{\mathcal{Q}_G}$ has a squarefree quadratic initial ideal if G is either a chordal graph or a ring graph (Proposition 1.4). In Section 2, we discuss the normality of the stable set polytopes. We prove that, for a simple graph G of stability number two, $\mathcal{Q}_G$ is normal if and only if the complement of G satisfies the "odd cycle condition" (Theorem 2.1). Using this criterion, we construct an infinite family of normal stable set polytopes without regular unimodular triangulations (Theorem 2.2). For general simple graphs, some necessary conditions for $\mathcal{Q}_G$ to be normal are also given. In Section 3, we study the set of generators and Gröbner bases of toric ideals of stable set polytopes. It is shown that, for a simple graph G of stability number two, the set of binomial generators of $I_{\mathcal{Q}_G}$ is described by the even closed walk of a graph (Theorem 3.2). If $\overline{G}$ is bipartite and if $I_{\mathcal{Q}_G}$ is generated by quadratic binomials, then $I_{\mathcal{Q}_G}$ has a quadratic Gröbner basis (Corollary 3.4). Finally, using the results on normality, generators, and Gröbner bases, we give an infinite family of nonnormal stable set polytopes whose toric ideal is generated by quadratic binomials and has no quadratic Gröbner bases (Theorem 2.2).

1. Fundamental properties of the stable set polytopes

In this section, we give some fundamental properties of $K[\mathcal{Q}_G]$ and $I_{\mathcal{Q}_G}$. In particular, a relation between the stable set polytopes and the edge polytopes is discussed. The *stability number* $\alpha(G)$ of a graph G is the cardinality of the largest stable set.

Example 1.1. If a simple graph G satisfies $\alpha(G) = 1$, i.e., G is a complete graph, then $I_{\mathcal{Q}_G} = \{0\}$, $K[\mathcal{Q}_G] = K[x_1t, \ldots, x_nt, t] \simeq K[y_1, \ldots, y_{n+1}]$, and $\mathcal{Q}_G$ is a simplex.

Example 1.2. Suppose that a simple graph G is not connected. Let $G_1, \ldots, G_s$ be connected components of G. Then, it is easy to see that $K[\mathcal{Q}_G]$ is isomorphic to the Segre product of $K[\mathcal{Q}_{G_1}], \ldots, K[\mathcal{Q}_{G_s}]$.

Thus, it is enough to study stable set polytopes of connected simple graphs G such that $\alpha(G) \geq 2$. The notion of toric fiber products [23] is a generalization of the Segre product. It is known [2] that we can apply the theory of toric fiber products to the toric rings of stable set polytopes. For $i = 1, 2$, let G_i be a simple graph on the vertex set V_i and the edge set E_i. If $V_1 \cap V_2$ is a clique of both G_1 and G_2, then we construct a new graph $G_1 \sharp G_2$ on the vertex set $V_1 \cup V_2$ and the edge set $E_1 \cup E_2$ which is called the *clique sum* of G_1 and G_2 along $V_1 \cap V_2$.

Proposition 1.3. *Let $G_1 \sharp G_2$ be the clique sum of simple graphs G_1 and G_2. Then, $I_{\mathcal{Q}_{G_1 \sharp G_2}}$ is a toric fiber product of $I_{\mathcal{Q}_{G_1}}$ and $I_{\mathcal{Q}_{G_2}}$. We can construct a set of binomial generators (or a Gröbner basis) of $I_{\mathcal{Q}_{G_1 \sharp G_2}}$ from that of $I_{\mathcal{Q}_{G_i}}$'s and some quadratic binomials. Moreover, $K[\mathcal{Q}_{G_1 \sharp G_2}]$ is normal if and only if both $K[\mathcal{Q}_{G_1}]$ and $K[\mathcal{Q}_{G_2}]$ are normal.*

Proof. This is a special case of [2, Proposition 5.1]. Note that $I_{\mathcal{Q}_{G_1 \cap G_2}} = \{0\}$. □

A simple graph G is called *chordal* if any induced cycle of G is of length 3. A *ring graph* is a graph whose block which is not a bridge or a vertex can be constructed from a cycle by successively adding cycles of length ≥ 3 using the edge sum construction. Ring graphs are introduced in [3, 4] and the toric ideals of cut polytopes of ring graphs are studied in [8].

Proposition 1.4. *Suppose that a simple graph G is either a chordal graph or a ring graph. Then, $I_{\mathcal{Q}_G}$ has a squarefree quadratic initial ideal.*

Proof. It is known that a graph G is chordal if and only if G is a clique sum of complete graphs. By the statement in Example 1.1 and Proposition 1.3, $I_{\mathcal{Q}_G}$ has a squarefree quadratic initial ideal if G is chordal.

Suppose that G is a ring graph. Then, G is a clique sum of trees and cycles. Since trees and cycles are almost bipartite, by [2, Theorem 8.1], the toric ideal $I_{\mathcal{Q}_H}$ has a squarefree quadratic initial ideal if H is either a tree or a cycle. Thus, by Proposition 1.3, $I_{\mathcal{Q}_G}$ has a squarefree quadratic initial ideal if G is a ring graph. □

A graph G is called *perfect* if the chromatic number of every induced subgraph of G equals the size of the largest clique of that subgraph. See, e.g., [1]. The following is known ([5, 15, 22]).

Proposition 1.5. *Let G be a simple graph. Then, $\mathcal{Q}_G$ is compressed if and only if G is perfect. In particular, if G is perfect, then $\mathcal{Q}_G$ is normal.*

For a graph G on the vertex set $[n]$, let $\overline{G}$ denote the complement of a graph G. An induced cycle of G of length > 3 is called *hole* of G. An induced cycle of $\overline{G}$ of length > 3 is called *antihole* of G. The first statement of Proposition 1.6 is called the *strong perfect graph theorem*.

Proposition 1.6. *Let G be a simple graph. Then, G is a perfect graph if and only if G has no odd holes and no odd antiholes. In particular, G is a perfect graph with $\alpha(G) = 2$ if and only if $\overline{G}$ is bipartite and not empty.*

For a graph G, let $\overline{G}^\star$ be the nonsimple graph on the vertex set $[n+1]$ whose edge (and loop) set is $E(\overline{G}) \cup \{\{i, n+1\} \ : \ i \in [n+1]\}$. The following lemma will play an important role when we study the stable set polytope $\mathcal{Q}_G$ of G.

Lemma 1.7. *Let G be a simple graph with $\alpha(G) = 2$. Then, we have $K[\mathcal{Q}_G] \simeq K[\mathcal{P}_{\overline{G}^\star}]$. Moreover, if $\overline{G}$ is bipartite, then there exists a bipartite graph H such that $K[\mathcal{Q}_G] \simeq K[\mathcal{P}_H]$.*

Proof. Let $\varphi : K[\mathcal{Q}_G] \to K[x_1, \ldots, x_{n+1}]$ be the injective ring homomorphism defined by $\varphi(x_1^{a_1} \cdots x_n^{a_n} t) = x_1^{a_1} \cdots x_n^{a_n} x_{n+1}^{2-(a_1+\cdots+a_n)}$. Then, $\varphi(t) = x_{n+1}^2$, $\varphi(x_i t) = x_i x_{n+1}$ for $i = 1, 2, \ldots, n$, and $\varphi(x_k x_\ell t) = x_k x_\ell$ for each stable set $\{k, \ell\}$ of G. Note that $\{k, \ell\}$ is a stable set of G if and only if $\{k, \ell\}$ is an edge of $\overline{G}$. Hence, the image of φ is $K[\mathcal{P}_{\overline{G}^\star}]$.

Suppose that $\overline{G}$ is bipartite. Then, $\overline{G}$ has no odd cycles. Hence, any odd cycle of $\overline{G}^\star$ has the vertex $n+1$. (Note that $\{n+1, n+1\}$ is an odd cycle of length

1.) Thus, in particular, any two odd cycles of $\overline{G}^\star$ has a common vertex. By the argument in [17, Proof of Proposition 5.5], there exists a bipartite graph H such that $K[\mathcal{Q}_G] \simeq K[\mathcal{P}_H]$. $\square$

The first application of Lemma 1.7 is as follows:

Proposition 1.8. *Let G be a simple graph. Then, the following conditions are equivalent:*

(i) *$\mathcal{Q}_G$ is unimodular;*
(ii) *$\overline{G}$ is bipartite.*

Moreover, if $\alpha(G) = 2$, then the conditions

(iii) *$\mathcal{Q}_G$ is compressed;*
(iv) *G is perfect*

are also equivalent to conditions (i) *and* (ii).

Proof. We may assume that G is not complete (i.e., $\overline{G}$ is not empty and $\alpha(G) \neq 1$). Let A be the matrix whose columns are vertices of $\mathcal{Q}_G$ and let

$$\widetilde{A} = \begin{pmatrix} & A & \\ 1 & \cdots & 1 \end{pmatrix} \text{ and } B = \begin{pmatrix} \mathbf{0} & \mathbf{e}_1 & \cdots & \mathbf{e}_n \\ 1 & 1 & \cdots & 1 \end{pmatrix}.$$

Then, B is a submatrix of $\widetilde{A}$. Since $|\det(B)| = 1$, it is known [20, p.70] that $\mathcal{Q}_G$ is unimodular if and only if the absolute value of any nonzero $(n+1)$-minor of the matrix $\widetilde{A}$ is 1.

Suppose that $\overline{G}$ is not bipartite. Then, $\overline{G}$ has an odd cycle $C = (i_1, \ldots, i_{2\ell+1})$. The the absolute value of the $(n+1)$-minor of $\widetilde{A}$ that corresponds to

$$\{\mathbf{e}_{i_k} + \mathbf{e}_{i_{k+1}} : 1 \leq k \leq 2\ell\} \cup \{\mathbf{e}_{i_1} + \mathbf{e}_{i_{2\ell+1}}, \mathbf{0}\} \cup \{\mathbf{e}_j : j \notin \{i_1, \ldots, i_{2\ell+1}\}\}$$

equals to 2. Hence, $\mathcal{Q}_G$ is not unimodular. Thus, we have (i) $\Rightarrow$ (ii).

Suppose that $\overline{G}$ is bipartite. By Lemma 1.7, there exists a bipartite graph H such that $K[\mathcal{Q}_G] \simeq K[\mathcal{P}_H]$. It is well-known that the edge polytope of a bipartite graph is unimodular. Thus, $\mathcal{P}_H$ is unimodular and hence we have (ii) $\Rightarrow$ (i).

Suppose that $\alpha(G) = 2$. By Proposition 1.5, conditions (iii) and (iv) are equivalent. In addition, by Proposition 1.6, conditions (ii) and (iv) are equivalent. $\square$

We close this section with the fundamental fact on stable set and edge polytopes.

Proposition 1.9. *Let G' be an induced subgraph of a graph G. Then,*

(i) *The edge polytope $\mathcal{P}_{G'}$ is a face of $\mathcal{P}_G$;*
(ii) *If G is a simple graph, then $\mathcal{Q}_{G'}$ is a face of $\mathcal{Q}_G$.*

By Proposition 1.9, a lot of the properties of $K[\mathcal{Q}_G]$ (resp. $K[\mathcal{P}_G]$) are inherited to $K[\mathcal{Q}_{G'}]$ (resp. $K[\mathcal{P}_{G'}]$); for example, the normality of the toric ring, the existence of a squarefree initial ideal, the existence of a quadratic Gröbner basis, the existence of the set of quadratic binomial generators of the toric ideal. See [10].

2. Normality of stable set polytopes

In this section, we study the normality of stable set polytopes. The normality of an edge polytope studied in [11, 19] will play an important role. If C_1 and C_2 are cycles in a graph G, then an edge $\{i, j\}$ of G is called a *bridge* of C_1 and C_2 if i is a vertex of $C_1 \setminus C_2$ and if j is a vertex of $C_2 \setminus C_1$. We say that a graph G satisfies the *odd cycle condition* if, for arbitrary two induced odd cycles C_1 and C_2 in G, either C_1 and C_2 have a common vertex or there exists a bridge of C_1 and C_2. For the sake of simpleness, assume that a graph H has at most one loop. Then, it is known [11, 19] that $\mathcal{P}_H$ is normal if and only if each connected component of H satisfies the odd cycle condition. By [11, Corollary 2.3] and Lemma 1.7, we have the following. (Note that $\overline{G}$ below is not necessarily connected.)

Theorem 2.1. *Let G be a simple graph with $\alpha(G) = 2$. Then, the following conditions are equivalent.*

(i) *$\mathcal{Q}_G$ is normal*;
(ii) *$\mathcal{Q}_G$ has a unimodular covering*;
(iii) *$\overline{G}$ satisfies the odd cycle condition, i.e., if two odd holes C_1 and C_2 in $\overline{G}$ have no common vertices, then there exists a bridge of C_1 and C_2 in $\overline{G}$.*

In particular, if $\mathcal{Q}_G$ is normal, then $\mathcal{P}_{\overline{G}}$ is normal.

Proof. Let G be a simple graph on the vertex set $[n]$ with $\alpha(G) = 2$. By Lemma 1.7, we have $K[\mathcal{Q}_G] \simeq K[\mathcal{P}_{\overline{G}^\star}]$. Hence, by [11, Corollary 2.3], conditions (i) and (ii) are equivalent, and they hold if and only if $\overline{G}^\star$ satisfies the odd cycle condition. (Note that $\overline{G}^\star$ is connected.) Since the vertex $n+1$ is incident with any vertices of $\overline{G}^\star$, it is easy to see that $\overline{G}^\star$ satisfies the odd cycle condition if and only if $\overline{G}$ satisfies the odd cycle condition. □

It is known [12] that there exists a graph G such that $\mathcal{P}_G$ is normal and that $I_{\mathcal{P}_G}$ has no squarefree initial ideals. In [9], infinite families of such edge polytopes are given. We can construct the stable set polytopes with the same properties. Let $G_1(p_1, \ldots, p_5)$ be a graph defined in [9, Theorem 3.10].

Theorem 2.2. *Let G be a graph whose complement is $G_1(p_1, \ldots, p_5)$ with $p_i \geq 2$ for $i = 1, \ldots, 5$. Then, $\mathcal{Q}_G$ is normal and $I_{\mathcal{Q}_G}$ has no squarefree initial ideals.*

Proof. Since $\overline{G}$ has no triangles, we have $\alpha(G) = 2$. Since $G_1(p_1, \ldots, p_5)$ satisfies the odd cycle condition, $\mathcal{Q}_G$ is normal by Theorem 2.1. On the other hand, $I_{\overline{G}}$ has no squarefree initial ideals. Since $\overline{G}$ is an induced subgraph of $\overline{G}^\star$, $I_{\mathcal{Q}_G}$ has no squarefree initial ideals by Lemma 1.7 and Proposition 1.9. □

It seems to be a challenging problem to characterize the normal steble set polytopes with large stability number. We give several necessary conditions. The following is a consequence of Proposition 1.9 and Theorem 2.1.

Proposition 2.3. *Let G be a simple graph. Suppose that $\mathcal{Q}_G$ is normal. Then, any two odd holes of $\overline{G}$ without common vertex have a bridge in $\overline{G}$.*

Proof. Suppose that two odd holes C_1, C_2 of $\overline{G}$ without common vertices have no bridges in $\overline{G}$. Let H be an induced subgraph of G whose vertex set is that of $C_1 \cup C_2$. Then, $\alpha(H) = 2$ and hence $\mathcal{Q}_H$ is not normal by Theorem 2.1. Thus, $\mathcal{Q}_G$ is not normal by Proposition 1.9. □

Similar conditions are required for antiholes of $\overline{G}$.

Proposition 2.4. *Let G be a simple graph. Suppose that $\mathcal{Q}_G$ is normal. Then, G satisfies all of the following conditions:*

(i) *Any two odd antiholes of $\overline{G}$ having no common vertices have a bridge in $\overline{G}$.*
(ii) *Any two odd antiholes of $\overline{G}$ of length ≥ 7 having exactly one common vertex have a bridge in $\overline{G}$.*
(iii) *Any odd hole and odd antihole of $\overline{G}$ having no common vertices have a bridge in $\overline{G}$.*

Proof. Let G be a graph on the vertex set $[n]$. Let $\mathcal{A} = \{(\rho(W), 1)^t : W \in S(G)\}$. It is known [20, Proposition 13.5] that $\mathcal{Q}_G$ is normal if and only if we have $\mathbb{Z}_{\geq 0}\mathcal{A} = \mathbb{Q}_{\geq 0}\mathcal{A} \cap \mathbb{Z}^{n+1}$.

(i) Let $C_1 = (i_1, \ldots, i_{2k+1})$ and $C_2 = (j_1, \ldots, j_{2\ell+1})$ be odd antiholes in $\overline{G}$ having no common vertices and no bridges in $\overline{G}$. By Proposition 1.9, we may assume that $\overline{G} = C_1 \cup C_2$. Then,

$$\begin{aligned} \sum_{W \in S(\overline{C_1}) \text{ and } |W|=k} (\rho(W), 1) &= (2k+1)\mathbf{e}_{n+1} + k\sum_{p=1}^{2k+1} \mathbf{e}_{i_p}, \\ \sum_{W \in S(\overline{C_2}) \text{ and } |W|=\ell} (\rho(W), 1) &= (2\ell+1)\mathbf{e}_{n+1} + \ell\sum_{q=1}^{2\ell+1} \mathbf{e}_{j_q}. \end{aligned}$$

Since $k, \ell \geq 2$, we have $k\ell - k - \ell \geq 0$. Hence,

$$\begin{aligned} & 5\mathbf{e}_{n+1} + \sum_{p=1}^{2k+1} \mathbf{e}_{i_p} + \sum_{q=1}^{2\ell+1} \mathbf{e}_{j_q} \\ = \ & \frac{1}{k}\left((2k+1)\mathbf{e}_{n+1} + k\sum_{p=1}^{2k+1} \mathbf{e}_{i_p}\right) + \frac{1}{\ell}\left((2\ell+1)\mathbf{e}_{n+1} + \ell\sum_{q=1}^{2\ell+1} \mathbf{e}_{j_q}\right) + \frac{k\ell - k - \ell}{k\ell}\mathbf{e}_{n+1} \end{aligned}$$

belongs to $\mathbb{Q}_{\geq 0}\mathcal{A} \cap \mathbb{Z}^{n+1}$. However, this vector is not in $\mathbb{Z}_{\geq 0}\mathcal{A}$ since $\max\{|W| : W \in S(\overline{C_1})\} = k$, $\max\{|W| : W \in S(\overline{C_2})\} = \ell$, and $\lceil (2k+1)/k \rceil + \lceil (2\ell+1)/\ell \rceil = 6 > 5$.

(ii) Let $C_1 = (i_1, \ldots, i_{2k+1})$ and $C_2 = (j_1, \ldots, j_{2\ell+1})$ be odd antiholes in $\overline{G}$ of length ≥ 7 having exactly one common vertex $i_1 = j_1$ and no bridges in $\overline{G}$. By Proposition 1.9, we may assume that $\overline{G} = C_1 \cup C_2$. Let

$$\begin{aligned} S_1 &= \{W \in S(\overline{C_1}) : |W| = k \text{ and either } i_1 \in W \text{ or } \{i_2, i_{2k+1}\} \subset W\}, \\ S_2 &= \{W \in S(\overline{C_2}) : |W| = \ell \text{ and either } j_1 \in W \text{ or } \{j_2, j_{2\ell+1}\} \subset W\}. \end{aligned}$$

Then,

$$\begin{aligned}\sum_{W\in S_1}(\rho(W),1) &= (2k-1)\mathbf{e}_{n+1}+k\mathbf{e}_{i_1}+(k-1)\sum_{p=2}^{2k+1}\mathbf{e}_{i_p},\\ \sum_{W\in S_2}(\rho(W),1) &= (2\ell-1)\mathbf{e}_{n+1}+\ell\mathbf{e}_{i_1}+(\ell-1)\sum_{q=2}^{2\ell+1}\mathbf{e}_{j_q}.\end{aligned}$$

Since $k,\ell\geq 3$, we have $0<1/(k-1)+1/(\ell-1)\leq 1$. Hence,

$$\begin{aligned}\alpha &:= 5\mathbf{e}_{n+1}+3\mathbf{e}_{i_1}+\sum_{p=2}^{2k+1}\mathbf{e}_{i_p}+\sum_{q=2}^{2\ell+1}\mathbf{e}_{j_q}\\ &= \frac{1}{k-1}\left((2k-1)\mathbf{e}_{n+1}+k\mathbf{e}_{i_1}+(k-1)\sum_{p=2}^{2k+1}\mathbf{e}_{i_p}\right)\\ &\quad+\frac{1}{\ell-1}\left((2\ell-1)\mathbf{e}_{n+1}+\ell\mathbf{e}_{i_1}+(\ell-1)\sum_{q=2}^{2\ell+1}\mathbf{e}_{j_q}\right)+\left(1-\frac{1}{k-1}-\frac{1}{\ell-1}\right)(\mathbf{e}_{i_1}+\mathbf{e}_{n+1})\end{aligned}$$

belongs to $\mathbb{Q}_{\geq 0}\mathcal{A}\cap\mathbb{Z}^{n+1}$. Suppose that $\alpha=(\rho(W_1),1)+\cdots+(\rho(W_5),1)$ for $W_i\in S(\overline{G})$. Then, each W_i belongs to either $S(\overline{C_1})$ or $S(\overline{C_2})$. Since $\max\{|W|:W\in S(\overline{C_1})\}=k$, $\max\{|W|:W\in S(\overline{C_2})\}=\ell$, we have

$$\begin{aligned}|\{W_1,\ldots,W_5\}\cap S(\overline{C_1})| &\geq 2,\\ |\{W_1,\ldots,W_5\}\cap S(\overline{C_2})| &\geq 2.\end{aligned}$$

We may assume that $W_1,W_2\in S(\overline{C_1})$ and $W_3,W_4,W_5\in S(\overline{C_2})$. It then follows that $\rho(W_1)+\rho(W_2)=\sum_{p=2}^{2k+1}\mathbf{e}_{i_p}$, and hence $\rho(W_3)+\rho(W_4)+\rho(W_5)=3\mathbf{e}_{i_1}+\sum_{q=2}^{2\ell+1}\mathbf{e}_{j_q}$. Hence $i_1\in W_3\cap W_4\cap W_5$. Thus, $i_2,i_{2\ell+1}\notin W_3,W_4,W_5$, a contradiction. Therefore, we have $\alpha\notin\mathbb{Z}_{\geq 0}\mathcal{A}$.

(iii) Let $C_1=(i_1,\ldots,i_{2k+1})$ be an odd hole and $C_2=(j_1,\ldots,j_{2\ell+1})$ an odd antihole in $\overline{G}$ having no common vertices. By Proposition 1.9, we may assume that $\overline{G}=C_1\cup C_2$. Then,

$$\begin{aligned}\sum_{W\in S(\overline{C_1})\text{ and }|W|=2}(\rho(W),1) &= (2k+1)\mathbf{e}_{n+1}+2\sum_{p=1}^{2k+1}\mathbf{e}_{i_p},\\ \sum_{W\in S(\overline{C_2})\text{ and }|W|=\ell}(\rho(W),1) &= (2\ell+1)\mathbf{e}_{n+1}+\ell\sum_{q=1}^{2\ell+1}\mathbf{e}_{j_q}.\end{aligned}$$

Hence,

$$\begin{aligned}&(k+3)\mathbf{e}_{n+1}+\sum_{p=1}^{2k+1}\mathbf{e}_{i_p}+\sum_{q=1}^{2\ell+1}\mathbf{e}_{j_q}\\ =\ &\frac{1}{2}\left((2k+1)\mathbf{e}_{n+1}+2\sum_{p=1}^{2k+1}\mathbf{e}_{i_p}\right)+\frac{1}{\ell}\left((2\ell+1)\mathbf{e}_{n+1}+\ell\sum_{q=1}^{2\ell+1}\mathbf{e}_{j_q}\right)+\frac{\ell-2}{2\ell}\mathbf{e}_{n+1}\end{aligned}$$

belongs to $\mathbb{Q}_{\geq 0}\mathcal{A}\cap\mathbb{Z}^{n+1}$. However, this vector is not in $\mathbb{Z}_{\geq 0}\mathcal{A}$ since $\max\{|W| : W \in S(\overline{C_1})\} = 2$, $\max\{|W| : W \in S(\overline{C_2})\} = \ell$, and $\lceil(2k+1)/2\rceil + \lceil(2\ell+1)/\ell\rceil = k+4 > k+3$. □

Unfortunately, the above conditions are not sufficient to be normal in general. For example, if the length of the two odd antiholes of $\overline{G}$ without common vertices are long, then a lot of bridges in $\overline{G}$ seem to be needed.

3. Generators and Gröbner bases of $I_{\mathcal{Q}_G}$

For a toric ideal I, let $\mu(I)$ be the maximum degree of binomials in a minimal set of binomial generators of I. If $I = \{0\}$, then we set $\mu(I) = 0$. In this section, we study $\mu(I_{\mathcal{Q}_G})$ by using results on the toric ideals of edge polytopes.

Let G be a graph on the vertex set $[n]$ allowing loops and having no multiple edges. Let $E(G) = \{e_1, \ldots, e_m\}$ be a set of all edges and loops of G. The toric ideal $I_{\mathcal{P}_G}$ is the kernel of a homomorphism $\pi : K[y_1, \ldots, y_m] \to K[x_1, \ldots, x_n]$ defined by $\pi(y_i) = x_k x_\ell$ where $e_i = \{k, \ell\}$. A walk of length q of G connecting $v_1 \in [n]$ and $v_{q+1} \in [n]$ is a finite sequence of the form

$$\Gamma = (\{v_1, v_2\}, \{v_2, v_3\}, \ldots, \{v_q, v_{q+1}\}) \tag{1}$$

with each $\{v_k, v_{k+1}\} \in E(G)$. An *even* (resp. *odd*) *walk* is a walk of even (resp. odd) length. A walk Γ of the form (1) is called *closed* if $v_{q+1} = v_1$. Given an even closed walk $\Gamma = (e_{i_1}, e_{i_2}, \ldots, e_{i_{2q}})$ of G, we write f_Γ for the binomial

$$f_\Gamma = \prod_{k=1}^{q} y_{i_{2k-1}} - \prod_{k=1}^{q} y_{i_{2k}} \in I_{\mathcal{P}_G}.$$

We regard a loop as an odd cycle of length 1. It is known ([13, 20, 24]) that

Proposition 3.1. *Let G be a graph having at most one loop. Then, $I_{\mathcal{P}_G}$ is generated by all the binomials f_Γ, where Γ is an even closed walk of G. In particular, $I_{\mathcal{P}_G} = (0)$ if and only if each connected component of G has at most one cycle and the cycle is odd.*

The following theorem implies that the set of binomial generators of $I_{\mathcal{Q}_G}$ is also characterized by the graph theoretical terminology if $\alpha(G) = 2$.

Theorem 3.2. *Let G be a simple graph with $\alpha(G) = 2$. Then, $I_{\mathcal{Q}_G} = I_{\mathcal{P}_{\overline{G}}} + J$ where J is an ideal generated by quadratic binomials f_Γ where Γ is an even closed walk of $\overline{G}^\star$ that satisfies one of the following:*

(i) *$\Gamma = (\{i, j\}, \{j, k\}, \{k, n+1\}, \{n+1, i\})$ is a cycle where $\{i, j\}$, $\{j, k\} \in E(\overline{G})$;*

(ii) $\Gamma = (\{i,j\},\{j,n+1\},\{n+1,n+1\},\{n+1,i\})$ *where* $\{i,j\} \in E(\overline{G})$.

In particular, $\mu(I_{\mathcal{Q}_G}) = \max\{\mu(I_{\mathcal{P}_{\overline{G}}}), 2\}$.

Proof. Let Γ be an even closed walk of $\overline{G}^\star$. It is enough to show that $f_\Gamma \in I_{\mathcal{P}_{\overline{G}^\star}}$ belongs to $I_{\mathcal{P}_{\overline{G}}} + J$. Suppose that f_Γ does not belong to $I_{\mathcal{P}_{\overline{G}}} + J$. Then, the vertex $n+1$ belongs to Γ. We may assume that the degree of f_Γ is minimum among such binomials. Then, f_Γ is irreducible. Let

$$\Gamma = (\{n+1,p\},\{p,q\},\{q,r\},\Gamma').$$

Since f_Γ is irreducible, it follows that p,q,r are distinct each other and that $q \neq n+1$. Then, f_Γ is generated by f_{Γ_1} and f_{Γ_2} where $\Gamma_1 = (\{n+1,p\},\{p,q\},\{q,r\},\{r,n+1\})$ and $\Gamma_2 = (\{n+1,r\},\Gamma')$. Since $\deg f_{\Gamma_2} < \deg f_\Gamma$, the binomial f_{Γ_2} belongs to $I_{\mathcal{P}_{\overline{G}}} + J$ by the assumption on $\deg f_\Gamma$. Moreover, Γ_1 satisfies one of the conditions (i) or (ii) and hence $f_{\Gamma_1} \in J$. Thus, we have $f_\Gamma \in \langle f_{\Gamma_1}, f_{\Gamma_2}\rangle \subset I_{\mathcal{P}_{\overline{G}}} + J$, a contradiction. □

Remark 3.3. A graph theoretical characterization of a *simple* graph G such that $\mu(I_{\mathcal{P}_G}) \leq 2$ is given in [13]. By Theorem 3.2, the characterization [13, Theorem 1.2] gives a graph theoretical characterization of a simple graph G such that $\alpha(G) = 2$ and $\mu(I_{\mathcal{Q}_G}) \leq 2$.

It is known [2, Theorem 7.4] that, if the complement of a graph G is an even cycle of length $2k$, then we have $\mu(I_{\mathcal{Q}_G}) = k$. By Theorem 3.2, we can generalize this result for a graph whose complement is an arbitrary bipartite graph.

Corollary 3.4. *Let G be a simple graph such that $\overline{G}$ is bipartite. Then, we have*

$$\mu(I_{\mathcal{Q}_G}) = \begin{cases} 0 & \text{if } \overline{G} \text{ is empty (i.e., } G \text{ is complete),} \\ k & \text{if } \overline{G} \text{ has a cycle,} \\ 2 & \text{otherwise,} \end{cases}$$

where $2k$ is the maximum length of induced cycles of $\overline{G}$. Moreover, the following conditions are equivalent:

(i) $\mu(I_{\mathcal{Q}_G}) \leq 2$, *i.e., $I_{\mathcal{Q}_G}$ is generated by quadratic binomials;*
(ii) $K[\mathcal{Q}_G]$ *is Koszul;*
(iii) $I_{\mathcal{Q}_G}$ *has a quadratic Gröbner basis;*
(iv) *The length of any induced cycle of $\overline{G}$ is* 4.

Proof. Let G be a simple graph such that $\overline{G}$ is bipartite. Then, $\alpha(G) = 2$. Since $\overline{G}$ is bipartite, it is known (see, e.g., [18]) that, $I_{\mathcal{P}_{\overline{G}}}$ is generated by f_Γ where Γ is an induced even cycle of $\overline{G}$. Note that $\deg f_\Gamma = k$ if the length of Γ is $2k$. Hence, by Theorem 3.2, we have a formula for $\mu(I_{\mathcal{Q}_G})$.

By this formula, it follows that (i) and (iv) are equivalent. Moreover, (iii) ⇒ (ii) ⇒ (i) holds in general. By Lemma 1.7, there exists a bipartite graph H such that $K[\mathcal{Q}_G] \simeq K[\mathcal{P}_H]$. By the theorem in [14], $I_{\mathcal{P}_H}$ has a quadratic Gröbner basis if and only if $\mu(I_{\mathcal{P}_H}) \leq 2$. Thus, we have (i) ⇒ (iii). □

If $\overline{G}$ is not bipartite, then condition (i) and (iii) in Corollary 3.4 are not equivalent. In order to construct an infinite family of counterexamples, the following proposition is important. (Proof is essentially given in [13, Proof of Proposition 1.6].)

Proposition 3.5. *Let P be a $(0,1)$-polytope. If I_P has a quadratic Gröbner basis, then the initial ideal is generated by squarefree monomials and hence P is normal.*

Theorem 3.6. *Let G be a simple graph such that $\overline{G}$ consists of two odd holes without common vertices. Then, $\alpha(G) = 2$ and*

(a) *$I_{\mathcal{Q}_G}$ is generated by quadratic binomials;*
(b) *$I_{\mathcal{Q}_G}$ has no quadratic Gröbner bases;*
(c) *$\mathcal{Q}_G$ is not normal.*

Proof. Since $\overline{G}$ has no triangles, we have $\alpha(G) = 2$. By Proposition 3.1 and Theorem 3.2, it follows that $I_{\overline{G}} = \{0\}$ and $I_{\mathcal{Q}_G}$ is generated by quadratic binomials. On the other hand, by Theorem 2.1, $\mathcal{Q}_G$ is not normal. Thus, by Proposition 3.5, $I_{\mathcal{Q}_G}$ has no quadratic Gröbner bases. □

The graphs in Theorem 3.6 are not strongly Koszul by the result in [7]. However, we do not know whether they are Koszul or not in general.

Remark 3.7. It is known [13, Theorem 1.2] that, if G is a simple connected graph and $I_{\mathcal{P}_G}$ is generated by quadratic binomials, then G satisfies the odd cycle condition and hence $\mathcal{P}_G$ is normal.

It seems to be a challenging problem to characterize the graphs G such that $\alpha(G) > 2$ and $\mu(I_{\mathcal{Q}_G}) \leq 2$. The following is a consequence of Proposition 1.9, Theorem 3.2 and [13, Theorem 1.2].

Proposition 3.8. *Let G be a simple graph. If $I_{\mathcal{Q}_G}$ is generated by quadratic binomials, then $\overline{G}$ satisfies the following conditions:*

(i) *Any even cycle of $\overline{G}$ of length ≥ 6 has a chord;*
(ii) *Any two odd holes of $\overline{G}$ having exactly one common vertex have a bridge;*
(iii) *Any two odd holes of $\overline{G}$ having no common vertex have at least two bridges.*

Proof. Suppose that $\overline{G}$ does not satisfy one of the conditions above. If $\overline{G}$ does not satisfy condition (i), then let H be an induced subgraph of G whose vertex set is that of the even cycle. If $\overline{G}$ does not satisfy condition either (ii) or (iii), then let H be an induced subgraph of G whose vertex set is that of two odd holes. Then, $\alpha(H) = 2$ and hence $I_{\mathcal{Q}_H}$ is not generated by quadratic binomials by Theorem 3.2 and [13, Theorem 1.2]. Thus, it follows from Proposition 1.9 that $I_{\mathcal{Q}_G}$ is not generated by quadratic binomials. □

References

[1] R. Diestel, "Graph Theory," Fourth Edition, Graduate Texts in Mathematics **173**, Springer, 2010.
[2] A. Engström and P. Norén, Ideals of graph homomorphisms, *Ann. Comb.* **17** (2013), 71–103.
[3] I. Gitler, E. Reyes and R. H. Villarreal, Ring graphs and toric ideals, *Electron. Notes Discrete Math.*, **28** (2007), 393–400.
[4] I. Gitler, E. Reyes and R. H. Villarreal, Ring graphs and complete intersection toric ideals, *Discrete Math.* **310** (2010), 430–441.
[5] J. Gouveia, P. A. Parrilo and R. R. Thomas, Theta bodies for polynomial ideals, *SIAM J. Optim.* **20** (2010), 2097–2118.

[6] T. Hibi and N. Li, Chain polytopes and algebras with straightening laws, *Acta Math. Vietnam.* **40** (2015) 447–452.
[7] K. Matsuda, Strong Koszulness of toric rings associated with stable set polytopes of trivially perfect graphs, *J. Algebra Appl.* **13** (2014), 1350138 [11 pages].
[8] U. Nagel and S. Petrović, Properties of cut ideals associated to ring graphs, *J. Commut. Algebra* **1** (2009) 547–565.
[9] H. Ohsugi, Toric ideals and an infinite family of normal $(0, 1)$-polytopes without unimodular regular triangulations, *Discrete Comput. Geom.* **27** (2002), 551–565.
[10] H. Ohsugi, J. Herzog and T. Hibi, Combinatorial pure subrings, *Osaka J. Math.* **37** (2000), 745–757.
[11] H. Ohsugi and T. Hibi, Normal polytopes arising from finite graphs, *J. Algebra* **207** (1998), 409–426.
[12] H. Ohsugi and T. Hibi, A normal $(0, 1)$-polytope none of whose regular triangulations is unimodular, *Discrete Comput. Geom.* **21** (1999), 201–204.
[13] H. Ohsugi and T. Hibi, Toric ideals generated by quadratic binomials, *J. Algebra* **218** (1999), 509–527.
[14] H. Ohsugi and T. Hibi, Koszul bipartite graphs, *Adv. Appl. Math.* **22** (1999), 25–28.
[15] H. Ohsugi and T. Hibi, Convex polytopes all of whose reverse lexicographic initial ideals are squarefree, *Proc. Amer. Math. Soc.*, **129** (2001), No.9, 2541–2546.
[16] H. Ohsugi and T. Hibi, Special simplices and Gorenstein toric rings, *J. Combin. Theory Ser. A* **113** (2006), Issue 4, 718–725.
[17] H. Ohsugi and T. Hibi, Centrally symmetric configurations of integer matrices, *Nagoya Math. J.* **216** (2014), 153–170.
[18] A. Simis, On the Jacobian module associated to a graph, *Proc. Amer. Math. Soc.*, **126** No. 4 (1998), 989–997.
[19] A. Simis, W. V. Vasconcelos and R. H. Villarreal, The integral closure of subrings associated to graphs, *J. Algebra* **199** (1998), 281–289.
[20] B. Sturmfels, "Gröbner bases and convex polytopes," Amer. Math. Soc., Providence, RI, 1996.
[21] B. Sturmfels, Four counterexamples in combinatorial algebraic geometry, *J. Algebra* **230** (2000) 282–294.
[22] S. Sullivant, Compressed polytopes and statistical disclosure limitation, *Tohoku Math. J.* **58** (2006), 433–445.
[23] S. Sullivant, Toric fiber products, *J. Algebra* **316** (2007), 560–577.
[24] R. Villarreal, Rees algebras of edge ideals, *Comm. Algebra*, **23** (1995), 3513–3524.

Kazunori Matsuda, Department of Pure and Applied Mathematics, Graduate School of Information Science and Technology, Osaka University, Suita, Osaka 565-0871, Japan
E-mail address: kaz-matsuda@ist.osaka-u.ac.jp

Hidefumi Ohsugi, Department of Mathematical Sciences, School of Science and Technology, Kwansei Gakuin University, Sanda, Hyogo, 669-1337, Japan
E-mail address: ohsugi@kwansei.ac.jp

Kazuki Shibata, Department of Mathematics, College of Science, Rikkyo University, Toshima-ku, Tokyo 171-8501, Japan
E-mail address: k-shibata@rikkyo.ac.jp